

NALOXONE PROVIDERS

Pharmacies

Walgreens Pharmacy • CVS Pharmacy
Smith's Food and Drug

Community-based Organizations

Northern Nevada HOPES

580 W 5th St Reno, NV 89503
775.786.4673

TracB-Exchange

6114 W Charleston Blvd Las Vegas, NV 89146
702.840.6693

Treatment Providers

Center for Behavioral Health

3050 E Desert Inn #116 Las Vegas, NV 89121
702.796.0660

2290 McDaniel St, Suite 1C North Las Vegas, NV 89030
702.399.1600

3470 W Cheyenne Rd, Suite 400 North Las Vegas, NV 89032
702.636.0085

1311 S Casino Center Blvd Las Vegas, NV 89104
702.382.6262

160 Hubbard Way, Suite A Reno, NV 89502
775.829.4472

Life Change Center

1201 N Stewart St Carson City, NV 89701
775.350.7250

1755 Sullivan Ln Sparks, NV 89431
775.355.7734

Vitality Unlimited

215 Bluffs Ave, Suite 200 Elko, NV 89801
800.242.8327

The average cash pay price range for
injectable naloxone (2 pack) in
Nevada is \$20-\$50

Injectable naloxone is covered under NV Medicaid
(with quantity limits)

For additional education and materials, please visit
prescribe365.nv.gov

Opioid Safety

and how to use naloxone



prescribe **365**
prescribe365.nv.gov



What is an opioid and how can you overdose?

Opioids include illicit drugs such as heroin or prescription medications used to treat pain such as oxycodone, hydrocodone, and fentanyl. An opioid overdose can happen when you take too many opioids at once or mix them with other drugs and your breathing slows or stops.



To avoid an accidental opioid overdose:

- Avoid mixing your opioids with alcohol, benzodiazepines (Xanax, Ativan, Klonopin, Valium), or medicines that make you sleepy.
- Be extra careful if you miss or change doses, feel ill, or start new medications.



How to identify an opioid overdose:

Look for these common signs:

- The person won't wake up or respond to voice or touch
- Breathing slows or even stops
- Lips and fingernails turn blue or gray
- Skin gets pale, clammy



In case of overdose:


- 1) Call 911 and give naloxone.
- 2) Do rescue breathing or chest compressions
 - Check for clear airway. If blocked, roll on side and use finger sweep to clear.
 - Roll onto back, tilt head back and pinch nose.
 - Give 2 regular breaths.
 - If still not breathing, give 1 breath every 5 seconds.
 - Continue until person revives or help arrives.
 - Once they start breathing, put them in the recovery position. Remember to keep breathing for them.
- 3) Place into Recovery Position:
 - Roll the person on their side, with one arm under their head.
 - Pull the top knee and arm over their body to keep them on their side. This will help prevent them from choking if they vomit. After naloxone, stay with a person for at least 3 hours or until help arrives.

How to administer injectable naloxone:

Step 1: Remove cap from naloxone vial and uncover the needle.

Step 2: Turn vial upside down and insert needle through rubber plug. Pull back on Plunger and draw 1ml (1cc) of liquid. Get rid of excess air.

Step 3: Inject into an upper arm, thigh muscle, or buttocks and empty the syringe.



Step 4: If no or minimal reaction in 2-3 minutes, give a second dose.

Nevada's Good Samaritan Law offers protection from arrest to individuals who seek medical assistance for a person experiencing a drug or alcohol overdose.

Protection is not offered from arrest for charges related to possession with intent to sell/distribute, child neglect/ endangerment or for individuals with active warrants for their arrest.

**Now that you have naloxone . . .
Tell someone where it is and how to use it.**