



References

Part I

- Babor & Higgins-Biddle (2001). *Brief intervention for hazardous and harmful drinking: A Manual for Use in Primary Care*. World Health Organization: Department of Mental Health and Substance Dependence.
- Berg, A.O. & Allan, J.D. (2001). Introducing the third U.S. Preventive Services Task Force. *American Journal of Preventive Medicine*, 20, 3–4.
- Bertholet, N., Daeppen, J.B., Wietlisbach, V., Fleming, M., & Burnand, B. (2005). Reduction of alcohol consumption by brief alcohol intervention in primary care: systematic review and meta-analysis. *Archives of Internal Medicine*, 165(9), 986-995.
- Coffield, F., Moseley, D., Hall, E., & Ecclestone, K. (2001). *Learning styles and pedagogy in post-16 learning: A systematic and critical review*. London: Learning and Skills Research Centre.
- Columbia Center for Addiction and Substance Abuse (2000). *Missed Opportunity: National Survey of Primary Care Physicians and Patients on Substance Abuse*. Survey Research Laboratory, University of Illinois at Chicago.
- Ferrer, R. L., Mody-Bailey, P., Jaén, C. R., Gott, S., & Araujo, S. (2009). A medical assistant–based program to promote healthy behaviors in primary care. *Annals of Family Medicine*, 7(6), 504–512.
- Fiellin, D. A., Reid, M. C., & O’Connor, P. G. (2000). Screening for alcohol problems in primary care: A systematic review. *Archives of Internal Medicine*, 160(13), 1977–1989.
- Friedmann, P., McCullough, D., Chin, M., Saitz, R. (2000). Screening and intervention for alcohol problems: A national survey of primary care physicians and psychiatrists. *Journal of General Internal Medicine*, 15(2), 84-91. http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1495340/pdf/jgi_03379.pdf
- Gentilello LM, Ebel BE, Wickizer TM, Salkever DS, Rivara FP. (2005). Alcohol interventions for trauma patients treated in emergency departments and hospitals: A cost benefit analysis. *Annals of Surgery*, 241, 541–550.
- Grumbach, K. & Bodenheimer, T. (2004). Can health care teams improve primary care practice? *Journal of the American Medical Association*, 291(10), 1246–1251.
- Hagemeier, N. E., Alamian, A., Murawski, M. M., Pack, R. P. (2015). Factors associated with provision of addiction treatment information by community pharmacists. *Journal of Substance Abuse Treatment*, 52, 67–72.
- Hasin, D.S., O’Brien, C.P., Auriacombe, M., Borges, G., Bucholz, K., Budney, A., Compton, W.M., Crowley, T., Ling, W., Petry, N.M., Schuckit, M., & Grant, B. F. (2013). DSM-5 criteria for substance use disorders: Recommendations and rationale. *American Journal of Psychiatry*, 170(8), 834-851.

- Hingson, R. & Compton, W. M. (2014). Screening and Brief Intervention and Referral to Treatment for Drug Use in Primary Care - Back to the Drawing Board. *JAMA*, 312(5). <http://www.sa4docs.org/wp-content/uploads/2014/08/Hingson-Compton-SBIRT-for-drug-use-in-primary-care-back-to-drawing-board-JAMA-8-5-14.pdf>
- Hung, D. Y., Rundall, T. G., Crabtree, B. F., Tallia, A. F., Cohen, D. J., & Halpin, H. A. (2006). Influence of primary care practice and provider attributes on preventive service delivery. *American Journal of Preventive Medicine*, 30(5), 413–422.
- Institute of Medicine Division of Mental Health and Behavioral Medicine. (1990). *Broadening the base of treatment for alcohol problems*. Washington, DC: National Academy Press.
- Kelly, T.M., Donovan, J.E., Chung, T., Bukstein, O.G., & Cornelius, J.R. (2009). Brief screens for detecting alcohol use disorder among 18-20 year old young adults in emergency departments: Comparing AUDIT-C, CRAFFT, RAPS4-QF, FAST, RUFT-Cut, and DSM-IV 2-item scale. *Addictive Behaviors*, 34(8), 668-774.
- McNeely, J., Cleland, C.M., Strauss, S.M., Palamar, J.J., Rotrosen, J., Saitz, R. (2015). Validation of self-administered single-item screening questions (SISQs) for unhealthy alcohol and drug use in primary care patients. *Journal of General Internal Medicine*, 30(12), 1757-1764.
- McNeely, J., Wright, S., Matthews, A. G., Rotrosen, J., Shelley, D., Buchholz, M. P., Curro, F.A., (2013). Substance-use screening and interventions in dental practices: Survey of practice-based research network dentists regarding current practices, policies and barriers. *JADA* 144(6), 627-638. <http://jada.ada.org>
- Mesnick, J. B. (2015). A shot of truth: Myth busting excessive alcohol use. Available online at <http://www.hhs.gov/blog/2015/02/23/shot-truth-myth-busting-excessive-alcohol-use.html>.
- Miller, P.M., Thomas, S.E., & Mallin, R. (2006). Patient attitudes towards self-report and biomarker alcohol screening by primary care physicians. *Alcohol & Alcoholism*, 41(3), 306-310.
- Mokdad, A.H., Marks, J.S., Stroup, D.F., Gerberding, J.L. (2004). Actual causes of death in the United States, 2000. *JAMA*, 291(10), 1238-1246.
- Moyer, A. Finney, J.W., Swearingen, C.E., & Vergun, P. (2002). Brief interventions for alcohol problems: A meta-analytic review of controlled investigations in treatment-seeking and non-treatment-seeking populations. *Addiction*, 97(3), 279-292.
- National Institute of Alcohol Abuse and Alcoholism (www.NIAAA.nih.gov)
- Russell, M. (2004/2005). *Screening in general health care*. *Alcohol Research and Health*, 28, 17-22.
- Sacks, J.J., Gonzales, K.R., Bouchery, E.E., Tomedi, L.E., & Brewer, R.D. (2015). 2010 National and state costs of excessive alcohol consumption. *American Journal of Preventive Medicine*, 49, e73-79.
- Smith, P.C., Schmidt, S.M., Allensworth-Davies, D., & Saitz, R. (2009). Primary care validation of a single-question alcohol screening test. *Journal of General Internal Medicine*, 24(7),783-788.
- Solberg, L.I., Maciosek, M.V., & Edwards, N.M. (2008). Primary care intervention to reduce alcohol misuse: Ranking its health impact and cost effectiveness. *American Journal of Prevention Medicine*, 43(2), 143.
- Substance Abuse and Mental Health Services Administration. (2014). *Results from the 2013 National Survey on Drug Use and Health: Summary of National Findings, NSDUH Series H-48, HHS Publication No. (SMA) 14-4863*. Rockville, MD: Substance Abuse and Mental Health Services Administration.

- United States Preventative Services Task Force. *Final Recommendation Statement Alcohol Misuse: Screening and Behavioral Counseling Interventions in Primary Care*.
<http://www.uspreventiveservicestaskforce.org/Page/Document/RecommendationStatementFinal/alcohol-misuse-screening-and-behavioral-counseling-interventions-in-primary-care>
- Vinson, D., Kruse, R.L., & Seale, J.P. (2007). Simplifying alcohol assessment: Two questions to identify alcohol use disorders. *Alcoholism, Clinical and Experimental Research*, 31(8), 1392–1398.
- Whitby, L.G. (1974). Screening for disease: Definitions and criteria. *Lancet*, 2, 819–822.
- Whilock, E.P., Polen, M.R., Green C.A., Orleans, T., Klein, J., U.S. Preventive Services Task Force. (2004). Behavioral counseling interventions in primary care to reduce risky/harmful alcohol use by adults: A summary of the evidence for the U.S. Preventive Services Task Force. *Annals of Internal Medicine*, 140(7), 557–568.
- Wilson, J.M.G. & Jungner, G. (1968). *Principles and Practice of Screening for Disease*. WHO Public Paper 34. Geneva: World Health Bibliography Organization.
- Wilson, L., Kahan, M., Liu, E., Brewster, J.M., Sobell, M.B., Sobell, L.C. (2002). Physician behavior towards male and female problem drinkers: a controlled study using simulated patients. *Journal of Addictive Disorders*, 21, 87–99. <http://www.ncbi.nlm.nih.gov/pubmed/12095002>
- Yersin, B., Nicolet, J.F., Dercrey, H., Burnier, M., van Melle, G., Pécoud, A. (1995). Screening for excessive alcohol drinking: Comparative value of carbohydrate-deficient transferrin, gamma-glutamyltransferase, and mean corpuscular volume. *Archives of Internal Medicine*, 155(17), 1907-11. <http://www.ncbi.nlm.nih.gov/pubmed/7677558>

Video: www.sbirtoregon.org

Part II

DAY 1

- Hasin, D.S., O'Brien, C.P., Auriacombe, M., Borges, G., Bucholz, K., Budney, A., Compton, W.M., Crowley, T., Ling, W., Petry, N.M., Schuckit, M., & Grant, B.F. (2013). DSM-5 criteria for substance use disorders: Recommendations and rationale. *American Journal of Psychiatry*, 170(8), 834-851.
- Smith, P.C., Schmidt, S.M., Allensworth-Davies, D., & Saitz, R. (2009). Primary care validation of a single-question alcohol screening test. *Journal of General Internal Medicine*, 24(7), 783-788.

DAY 2

- Miller, W.R. (2004). *Toward a Theory of Motivational Interviewing*. Available at motivationalinterviewing.org.
- Miller, W. R. (2010). *Motivational therapy sessions*. Alexandria, VA: Alexander Street Press.
- Miller, W.R. & Baca, L.M. (1983). Two-year follow-up of bibliotherapy and therapist-directed controlled drinking training for problem drinkers. *Behavior Therapy*, 14(3), 441-448.
- Miller, W.R., Benefield, R.G., & Tonigan, J.S. (1993). Enhancing motivation for change in problem drinking: A controlled comparison of two therapist styles. *Journal of Consulting and Clinical Psychology*, 61(3), 455-461.
- Miller, W.R., Taylor, C.A., & West, J.C. (1980). Focused versus broad spectrum behavior therapy for problem drinkers. *Journal of Consulting and Clinical Psychology*, 48(5), 590–601.